

Rigorous numerics for dissipative PDEs

P. Zgliczyński

Jagiellonian University, Kraków, Poland

<http://www.ii.uj.edu.pl/~zgliczyn>

Outline of this talk

1. Model problem - Kuramoto-Sivashinsky Eq. and our results about the dynamics
2. Algorithm for rigorous integration of dissipative PDEs
3. Some data from the proofs
4. Conclusions, future work

A Model Problem - Kuramoto-Sivashinsky PDE

Consider the Kuramoto-Sivashinsky (KS) eq.

$$u_t = -\nu u_{xxxx} - u_{xx} + 2uu_x, \quad \nu > 0$$

where $(t, x) \in [0, \infty) \times \mathbf{R}$ subject to periodic and odd boundary conditions

$$\begin{aligned} u(t, 0) &= u(t, 2\pi) \\ u(t, -x) &= -u(t, x) \end{aligned}$$

For various values of ν a variety of dynamics,

fixed points,
periodic orbits,
heteroclinic orbits,
chaotic dynamics,

have been observed numerically.

Goal: A rigorous means of proving these numerical results.

A Model Problem - Kuramoto-Sivashinsky PDE, Fourier expansion

Fourier expansion is: $u(t, x) = \sum_{k=-\infty}^{\infty} b_k(t) e^{ikx}$

Substituting in **KS** and applying boundary conditions gives:

$$\dot{a}_k = k^2(1 - \nu k^2)a_k - k \sum_{n=1}^{k-1} a_n a_{k-n} + 2k \sum_{n=1}^{\infty} a_n a_{n+k}$$

where $b_k = ia_k$ and $k = 1, 2, 3, \dots$

Linearization: $\dot{a}_k = k^2(1 - \nu k^2)a_k$

- k -th mode is unstable for $k < \frac{1}{\sqrt{\nu}}$
- k -th mode is stable for $k > \frac{1}{\sqrt{\nu}}$
- the modes with $k \gg \frac{1}{\sqrt{\nu}}$ should be irrelevant for the dynamics

A Model Problem - Kuramoto-Sivashinsky PDE, known results

Known results:

- the existence of global attractor, the functions from attractor are analytic - **Fourier series converge at geometric rate** (Foias, Temam)
- the existence of finite dimensional inertial manifold (Foias, Nicolaenko, Sell, Temam, Rossa, Jolly) (**not of much use in rigorous numerics**)

No analytical results dynamics more complicated than fixed points bifurcating from zero solution

Our rigorous results for Kuramoto-Sivashinsky PDE

- the existence of multiple periodic orbits for various parameter values $\nu \approx 0.1215, 0.1212, 0.125, 0.032, 0.02991$, both stable and unstable orbits
- the existence of multiple fixed points for various values of ν and their bifurcations (joint with K. Mischaikow)
- the existence of attractive fixed points for various values of ν

Periodic point for KS-equation

$$\mu = 0.127$$

Symmetric attracting orbit

Theorem: Let $u_0(x) = \sum_{k=1}^{10} -2a_k \sin(kx)$, where a_k are given in table below. There exists a function $u^*(t, x)$, the classical solution of KS for $\nu = 0.127$, such that

$$\begin{aligned} \|u_0 - u^*(0, \cdot)\|_{L_2} &< 8.1 \cdot 10^{-4}, \\ \|u_0 - u^*(0, \cdot)\|_{C^0} &< 6.5 \cdot 10^{-4} \end{aligned}$$

such that u^* is periodic with respect to t .

$a_1 = 2.012088e - 01$	$a_2 = 1.289978$
$a_3 = 2.012152e - 01$	$a_4 = -3.778654e - 01$
$a_5 = -4.231056e - 02$	$a_6 = 4.316137e - 02$
$a_7 = 6.940373e - 03$	$a_8 = -4.156441e - 03$
$a_9 = -7.945097e - 04$	$a_{10} = 3.315994e - 04$

Proof uses Brouwer Thm. and rigorous integration of KS-PDE

Periodic point for KS-equation

$$\mu = 0.1215$$

non-symmetric attracting orbit past period doubling

Theorem: Let $u_0(x) = \sum_{k=1}^{13} -2a_k \sin(kx)$, where a_k are given in table below. There exists a function $u^*(t, x)$, the classical solution of KS for $\nu = 0.1215$, such that

$$\begin{aligned} \|u_0 - u^*(0, \cdot)\|_{L_2} &< 9.9 \cdot 10^{-5}, \\ \|u_0 - u^*(0, \cdot)\|_{C^0} &< 6.2 \cdot 10^{-5} \end{aligned}$$

such that u^* is periodic with respect to t .

$a_1 = 2.559310e - 01$	$a_2 = 1.096696$
$a_3 = 2.559302e - 01$	$a_4 = -3.079615e - 01$
$a_5 = -4.780276e - 02$	$a_6 = 3.002052e - 02$
$a_7 = 7.352633e - 03$	$a_8 = -2.530197e - 03$
$a_9 = -7.561938e - 04$	$a_{10} = 1.624861e - 04$
$a_{11} = 6.833008e - 05$	$a_{12} = -8.789182e - 06$
$a_{13} = -5.429523e - 06$	

Proof uses Brouwer Thm. and rigorous integration of KS-PDE

Periodic point for KS-equation

$$\mu = 0.1215$$

Symmetric unstable orbit, past period doubling

Theorem: Let $u_0(x) = \sum_{k=1}^{11} -2a_k \sin(kx)$, where a_k are given in table below. There exists a function $u^*(t, x)$, the classical solution of KS for $\nu = 0.1215$, such that

$$\begin{aligned} \|u_0 - u^*(0, \cdot)\|_{L_2} &< 1.27 \cdot 10^{-3}, \\ \|u_0 - u^*(0, \cdot)\|_{C^0} &< 8.26 \cdot 10^{-4} \end{aligned}$$

such that u^* is periodic with respect to t .

$a_1 = 2.450027e - 01$	$a_2 = 1.041500e + 00$
$a_3 = 2.449985e - 01$	$a_4 = -2.760754e - 01$
$a_5 = -4.371320e - 02$	$a_6 = 2.531380e - 02$
$a_7 = 6.345919e - 03$	$a_8 = -1.996779e - 03$
$a_9 = -6.177148e - 04$	$a_{10} = 1.184863e - 04$
$a_{11} = 5.269771e - 05$	

Proof uses Miranda Thm. and rigorous integration of KS-PDE, the orbit is apparently unstable

Periodic point for KS-equation

$$\mu = 0.032$$

symmetric attracting orbit, close to chaotic region

Theorem: Let $u_0(x) = \sum_{k=1}^{23} -2a_k \sin(kx)$, where a_k are given in table below. There exists a function $u^*(t, x)$, the classical solution of KS for $\nu = 0.032$, such that

$$\begin{aligned} \|u_0 - u^*(0, \cdot)\|_{L_2} &< 8.9 \cdot 10^{-4}, \\ \|u_0 - u^*(0, \cdot)\|_{C^0} &< 9.5 \cdot 10^{-4} \end{aligned}$$

such that u^* is periodic with respect to t .

$a_1 = 3.506682e - 01$	$a_2 = 2.522889e - 02$
$a_3 = 3.506665e - 01$	$a_4 = -2.276745e + 00$
$a_5 = -1.115325e + 00$	$a_6 = -3.693057e - 01$
$a_7 = 4.603873e - 01$	$a_8 = -4.604564e - 01$
$a_9 = -3.115024e - 01$	$a_{10} = -1.449674e - 01$
$a_{11} = 5.104894e - 02$	$a_{12} = -2.165916e - 02$
$a_{13} = -3.413293e - 02$	$a_{14} = -2.613508e - 02$
$a_{15} = 1.307623e - 03$	$a_{16} = 8.752424e - 05$
$a_{17} = -2.115586e - 03$	$a_{18} = -2.891477e - 03$
$a_{19} = -5.007345e - 04$	$a_{20} = 3.374289e - 05$
$a_{21} = -4.423567e - 05$	$a_{22} = -2.280484e - 04$
$a_{23} = -9.029570e - 05$	

The method of self-consistent bounds

H - Hilbert space,

e_1, e_2, \dots - an orthogonal basis in H

The corresponding projections are

$$p_m = P_m a := (a_1, a_2, \dots, a_m)$$

$$q_m = Q_m a := (a_{m+1}, a_{m+2}, \dots)$$

The problem:

$$\dot{a} = F(a) \quad (1)$$

F is not continuous, with dense domain in H .

$F_k \circ P_n$ is a C^1 -function for $n, k \in \mathbb{N}$

Later $F(a) = L(a) + N(a)$, L - linear, N - non-linear

e_1, e_2, \dots - eigenvectors of L - very helpful

The method:

Def. Fix m, M ($m \leq M$). A compact set $W \subset P_m(H)$ and a sequence of pairs $\{a_k^\pm \in \mathbf{R} \mid a_k^- < a_k^+, k \in \mathbf{Z}^+\}$ are *self-consistent a-priori bounds for F* if:

C1 For $k > M$, $a_k^- < 0 < a_k^+$.

C2 Let $\hat{a}_k := \max |a_k^\pm|$ and set $\hat{u} = \sum_{k=0}^{\infty} \hat{a}_k e_k$.
Then, $\hat{u} \in H$, $(\{\hat{a}_k\} \in l_2)$

C3 The function $u \mapsto F(u)$ is continuous on

$$W \oplus \prod_{k=m+1}^{\infty} [a_k^-, a_k^+] \subset H.$$

Moreover, if we define

$$\hat{f}_k = \max_{u \in W \oplus \prod_{k=m+1}^{\infty} [a_k^-, a_k^+]} |F_k(u)| \text{ and set}$$

$$\hat{f} = \sum \hat{f}_k e_k, \text{ then } \hat{f} \in H. (\{\hat{f}_k\} \in l_2)$$

Notation: $T = \prod_{k=m+1}^{\infty} [a_k^-, a_k^+]$ - Tail

ISOLATION for $n > m$

For $a \in W \oplus T$ and $k > m$ holds

$$a_k = a_k^+ \quad \Rightarrow \quad \dot{a}_k < 0$$

$$a_k = a_k^- \quad \Rightarrow \quad \dot{a}_k > 0$$



C1,C2,C3 - give convergence

C4 - gives a priori bounds

C1,C2,C3,C4 - easy to satisfy (later)

Finite dimensional rigorous computations in m first variables

Basic Differential Inclusion:

$$\dot{p} \in P_m F(p) + \Gamma_m, \quad p \in \mathbf{R}^m, \quad (2)$$

where $\Gamma_m = \{P_m F(p + q) - P_m F(p) \mid q \in T\}$

We say a multivalued map $p_I : [0, h] \rightarrow H$ is *upper attainable set (uas) map* for (2) if the following is true

- any C^1 function satisfying (2) and defined on the maximum interval of existence is defined on $[0, h]$
- if a C^1 -function $p : [0, h] \rightarrow X_m$ satisfies (2), then $p(t) \in p_I(t)$ for $t \in [0, h]$

Theorem: Assume $W \oplus T$ are self-consistent bounds for F . If $p_I : [0, t_1] \rightarrow X_m = P_m(H)$ is uas map for (2), such that $p_I([0, t_1]) \subset W$.

Then for any $q_0 \in T$, the problem $u' = F(u)$ (and all its Galerkin projections $u' = P_n F(u)$, $n > M$) has a solution $u(t) = (p(t), q(t))$ for $t \in [0, t_1]$, such that

$$p(t) \in p_I(t), \quad q(t) \in T, \quad \text{for } t \in [0, t_1]$$

Why it is a easy to find a good tail =
self-consistent bounds

$$u_t = Lu + N(u, Du, \dots, D^r u)$$

$x \in \mathbf{T}^n$ (periodic boundary conditions),
 L - linear, diagonal, N - polynomial

Fourier expansion $u(t) = \sum_{k \in \mathbf{Z}^n} a_k(t) e^{ik \cdot x}$

Lemma. Let $s > s_0$. If $|a_k| \leq C/|k|^s$, $|a_0| \leq C$,
then there exists $D = D(C, s)$

$$|N_k| \leq \frac{D}{|k|^{s-r}}, \quad |N_0| \leq D$$

Isolation. Assume $L(a)_k = -|k|^p a_k$, $p > r$.

Assume $|a_k| \leq \frac{C}{|k^s|}$, $|a_{k_0}| = \frac{C}{|k_0|^s}$, then

$$\begin{aligned} \frac{d|a_{k_0}|}{dt} &\leq -|k_0|^p |a_{k_0}| + |N_{k_0}(a)| \leq \\ &\quad -C|k_0|^{p-s} + D|k_0|^{r-s} \\ \frac{d|a_{k_0}|}{dt} &< 0, \quad |k_0| > M \end{aligned}$$

Rigorous integration for dissipative PDEs

$(x, y) \in X_m \oplus Y_m \subset H$ - Hilbert space,

$\dim X_m = m < \infty$, $\dim Y_m \leq \infty$

$P_m : H \rightarrow H$, projection onto X_m , $Q_m = I - P_m$

F - our PDE in some basis on H

$$x' = PF(x, y) \tag{3}$$

$$y' = QF(x, y) \tag{4}$$

Idea: Replace (3 - 4) by

$$x'(t) \in P_m F(x(t), Tail(t)) \tag{5}$$

$$y(t) \in Tail(t), \tag{6}$$

where $Tail(t)$ has *finite representation* and can be computed *in finite number of operations*. $Tail_k(t)$ should decay fast enough.

We want also that: $x(t) \oplus P_n Q_m Tail(t)$, for $n > M$, is a rigorous estimate to n -dimensional Galerkin projection of F , for the initial condition $x(0) \oplus P_n Q_m Tail(0)$

Integration of dissipative PDEs - II

$$x'(t) \in P_m F(x(t), Tail(t)) \quad (7)$$

$$y(t) \in Tail(t), \quad (8)$$

$x' = P_m F(x)$ - Galerkin projection, induces φ_m

One time step:

initial condition $Z \oplus Tail(0) \subset X_m \oplus Y_m$, $h > 0$

1 • find $W \oplus T[0, h]$ (**rough enclosure**)

$$P_m F(x, y) - P_m F(x, 0) \subset \Gamma, \quad x \in Z$$

$$\varphi_{m, \Gamma}([0, h], Z) \subset W$$

$$Tail([0, h]) \subset T[0, h].$$

2 • instead of (7) consider $x' \in P_m F(x, 0) + \Gamma$
- use algorithm for differential inclusions, to obtain $x(h)$ for $(x, y) \in Z \oplus Tail(0)$.

3 • compute $Tail(h)$.

Representation used for KS equation

We look for solutions in

$$W \oplus T = W \oplus \prod_{k=m+1}^{k \leq M} \oplus \prod_{k > M} \left[\frac{-C}{k^s}, \frac{C}{k^s} \right] \quad (9)$$

where $W \subset X_m$.

$$N_k(W \oplus T) \subset [N_k^-, N_k^+], \quad k = m + 1, \dots, M$$
$$N_k(W \oplus T) \subset \left[\frac{-D(W \oplus T)}{k^{s-2}}, \frac{D(W \oplus T)}{k^{s-2}} \right], \quad k > M$$

We solve (estimate rigorously) the solutions of the following system of differential inclusions

$$x' \in P_m F(x) + \Gamma, \quad x \in W \subset X_m$$
$$x'_k \in \lambda_k x_k + [N_k^-, N_k^+], \quad k = m + 1, \dots,$$

x_k for $k > M$ are given by a single formula.

Rigorous integration for ODEs and differential inclusions - basic principles

$x \in \mathbf{R}^n$, $f : \mathbf{R}^n \rightarrow \mathbf{R}^n - C^1$.

$$x' = f(x), \quad x(0) = x_0 \quad (\text{ODE})$$

induces $\varphi(t, x_0) \in \mathbf{R}^n$

One time step:

initial condition: $X_0 \subset \mathbf{R}^n$, $h > 0$ is a time step

1• find $W \subset \mathbf{R}^n$ (rough enclosure), such that $\varphi([0, h], X_0) \subset W$

2• apply the Taylor method to (ODE), evaluate the error term on W to obtain $X_1 \subset \mathbf{R}^n$, such that

$$\varphi(h, X_0) \subset X_1$$

Rigorous integration for ODEs - comments

- all computations are performed in **interval arithmetic**
- one should be very careful in the way how step **2** is executed, straightforward interval evaluation leads to *the wrapping effect*.
- we use *the Lohner algorithm*

Rigorous integration of differential inclusion

Differential inclusion : $\Gamma \subset \mathbf{R}^n$

$$x' \in f(x) + \Gamma, \quad x(0) = x_0$$

induces $\varphi_\Gamma(t, x_0) \subset \mathbf{R}^n$

One time step:

initial condition: $X_0 \subset \mathbf{R}^n$, $h > 0$ is a time step

1• compute X_1 , such that $\varphi(h, X_0) \subset X_1$

2• find $W_2 \subset \mathbf{R}^n$ (**rough enclosure**), such that

$$\varphi_\Gamma([0, h], X_0) \subset W_2, \quad x \in X_0$$

3• use Gronwall type lemma to find $\Delta \subset \mathbf{R}^n$,

$$\varphi_\Gamma(h, x) - \varphi(h, x) \in \Delta$$

This step requires $\frac{\partial f}{\partial x}(W_2)$

4•

$$\varphi_\Gamma(h, X_0) \subset X_1 + \Delta$$

Differential inclusions - Fundamental Lemma

For a fixed $y_c \in \mathbf{R}^{n_2}$ we compare the solutions of two ODEs

$$\begin{aligned}x_1' &= f(x_1, y_c), \\x_2' &= f(x_2, y_c) + (f(x_2, y(t)) - f(x_2, y_c)) \\x_1(t_0) &= x_2(t_0) = x_0\end{aligned}$$

where $y(t)$ is given (but unknown) function.

Lemma: Let:

$[W_y] \subset \mathbf{R}^{n_2}$, convex, $y([t_0, t_0 + h]) \subset [W_y]$.

$[W_1] \subset [W_2] \subset \mathbf{R}^{n_1}$ - convex and compact.

$x_1([t_0, t_0 + h]) \subset [W_1]$, $x_2([t_0, t_0 + h]) \subset [W_2]$ for any continuous function $y : [t_0, t_0 + h] \rightarrow [W_y]$.

Then the following inequality holds for $t \in [t_0, t_0 + h]$ and for $i = 1, \dots, n_1$

$$|x_{1,i}(t) - x_{2,i}(t)| \leq \left(\int_{t_0}^t e^{J(t-s)} C ds \right)_i, \quad (10)$$

where

$$[\delta] = \{f(x, y_c) - f(x, y) \mid x \in [W_1], y \in [W_y]\},$$

$$C_i \geq \sup \|\delta_i\|, \quad i = 1, \dots, n_1$$

$$J_{ij} \geq \sup \frac{\partial f_i}{\partial x_j}([W_2], [W_y]) \text{ if } i = j,$$

$$J_{ij} \geq \sup \left| \frac{\partial f_i}{\partial x_j}([W_2], [W_y]) \right| \text{ if } i \neq j.$$

Tail evolution

Our problem $a'_k = \lambda_k a_k + N_k(a)$,
 $\lambda_k \rightarrow -\infty$, for $|k| \rightarrow \infty$

$W, T([0, h])$ - the rough enclosure for $Z \oplus T(0)$
for $t \in [0, h]$

For $k > m$ we have

$$N_k^\pm = N_k^\pm(W, T([0, h]))$$
$$\lambda_k a_k + N_k^- < \frac{da_k}{dt} < \lambda_k a_k + N_k^+,$$

hence

$$b_k^\pm = \frac{N_k^\pm}{-\lambda_k}, \quad (11)$$

$$T(h)_k^\pm = \left(T(0)_k^\pm - b_k^\pm \right) e^{\lambda_k h} + b_k^\pm \quad (12)$$

It remains to put $T(h)$ for $k > M$ in the form

$$T(h)_k^\pm = \frac{\pm C(T(h))}{k^s(T(h))}$$

For $k > M$ we have

$$\begin{aligned}
 0 &< b_k^+ \leq \frac{C(b)}{k^{s(b)}} \\
 T(0)_k^+ &= \frac{C(T(0))}{k^{s(T(0))}} \\
 T(h)_k^+ &\leq T(0)_k^\pm e^{\lambda_k h} + b_k^\pm.
 \end{aligned}$$

$$T(h)_k^+ \leq \frac{C(T(0))}{k^{s(T(0))}} e^{\lambda_k h} + \frac{C(b)}{k^{s(b)}}.$$

Let

$$E = e^{h\lambda_{M+1}} (M+1)^{s(b)-s(T(0))}.$$

then (modulo some conditions on M, h)

$$e^{\lambda_k h} \leq \frac{E}{k^{s(b)-s(T(0))}}, \quad k > M$$

and finally we can set

$$T_k^\pm(h) = \pm \frac{C(T(0))E + C(b)}{k^{s(b)}}.$$

About the computations

- gnu C++
- interval arithmetic - from CAPD package developed in Krakow, Poland
- we use the Lohner algorithm to integrate differential inclusions

Some computation data

On 3GHz machine, Linux, gnu C++

- $\nu = 0.127$, $m = 10$, $M = 3 * m$, $h = 1e - 3$, $order = 4$, $T/2 \approx 1.12$, computation time around 10 sec
- $\nu = 0.1215$, $m = 13$, $M = 3 * m$, $h = 4e - 4$, $order = 6$, $T \approx 3.07$, computation time around 240 sec
- $\nu = 0.032$, $m = 23$, $M = 3 * m$, $h = 1.5e - 4$, $order = 5$, $T/2 \approx 0.41$, computation time around 300 sec

Conclusions

- rigorous numerics for dissipative PDEs is possible
- global existence and uniqueness theorems are not required, interesting solutions are constructed
- could be applied to (I hope): Ginzburg-Landau, Navier-Stokes in 2D and 3D

Future work

- prove chaos (symbolic dynamics) for KS $\nu \approx 0.029$ or $\nu \approx 0.1212$
- Construct an *rigorous C^1 -algorithm* for dissipative PDE.

This will make possible to **rigorously** apply a lot of dynamical system theory to dissipative PDEs.